


Quantile–quantile plot.

Unit exponential distribution vs. expected events (Cox–Snell residuals).

Should follow line through origin at 45 degrees (blue) if well fit.


Complete model:

Surv(time, status == 2) ~ age + edema + bili + protime + albumin



Observed vs. expected hazard.
Should follow line through origin at 45 degrees (blue) if well fit.

Complete model:
Surv(time, status == 2) ~ age + edema + bili + protime + albumin




Expected events vs. hazard based on sorted expected events
or Cox–Snell residuals vs. cumulative hazard of these residuals.
Should follow line through origin at 45 degrees (blue) if well fit.

Hazard = cumulative sum of [events/ no. at risk] (sorted by expected events)


Complete model:

Surv(time, status == 2) ~ age + edema + bili + protime + albumin




Time vs. hazard, per predictor.
If hazards proportional then curves should be constant multiples of a baseline.
Reference (black) line is 45 degrees.

Predictor: age




Time vs. hazard, per predictor.
If hazards proportional then curves should be constant multiples of a baseline.
Reference (black) line is 45 degrees.

Predictor: edema




Time vs. hazard, per predictor.
If hazards proportional then curves should be constant multiples of a baseline.
Reference (black) line is 45 degrees.

Predictor: bili




Time vs. hazard, per predictor.
If hazards proportional then curves should be constant multiples of a baseline.
Reference (black) line is 45 degrees.

Predictor: protime




Time vs. hazard, per predictor.
If hazards proportional then curves should be constant multiples of a baseline.
Reference (black) line is 45 degrees.

Predictor: albumin




Time vs. difference in log hazards, per predictor.
Should be constant over time.
If >0 (black line) shows survival advantage for reference group.

Reference: age = [26.3,45.8)




Time vs. difference in log hazards, per predictor.
Should be constant over time.
If >0 (black line) shows survival advantage for reference group.

Reference: edema = 0.0




Time vs. difference in log hazards, per predictor.
Should be constant over time.
If >0 (black line) shows survival advantage for reference group.

Reference: bili = [0.3, 1.0)




Time vs. difference in log hazards, per predictor.
Should be constant over time.
If >0 (black line) shows survival advantage for reference group.

Reference: protime = [9.0,10.3)




Time vs. difference in log hazards, per predictor.
Should be constant over time.
If >0 (black line) shows survival advantage for reference group.

Reference: albumin = [1.96,3.37)




Cumulative hazard vs. reference group. Should be linear plot through origin.
If convex (towards top left) shows ratio of hazards is increasing over time.
Reference line (black) is at 45 degrees.

Predictor: age




Cumulative hazard vs. reference group. Should be linear plot through origin.
If convex (towards top left) shows ratio of hazards is increasing over time.
Reference line (black) is at 45 degrees.

Predictor: edema




Cumulative hazard vs. reference group. Should be linear plot through origin.
If convex (towards top left) shows ratio of hazards is increasing over time.
Reference line (black) is at 45 degrees.

Predictor: bili




Cumulative hazard vs. reference group. Should be linear plot through origin.
If convex (towards top left) shows ratio of hazards is increasing over time.
Reference line (black) is at 45 degrees.

Predictor: protime




Cumulative hazard vs. reference group. Should be linear plot through origin.
If convex (towards top left) shows ratio of hazards is increasing over time.
Reference line (black) is at 45 degrees.


Predictor: albumin



Poisson approach to determine non-linearity.
Predictor (with smoothing spline) vs. residuals from GAM plot.
If linear should be horizontal line with intercept=0 (blue).




Poisson approach to determine non-linearity.
Predictor (with smoothing spline) vs. residuals from GAM plot.
If linear should be horizontal line with intercept=0 (blue).




Poisson approach to determine non-linearity.
Predictor (with smoothing spline) vs. residuals from GAM plot.
If linear should be horizontal line with intercept=0 (blue).

Predictor: s(protime, 3)




Poisson approach to determine non-linearity.
Predictor (with smoothing spline) vs. residuals from GAM plot.
If linear should be horizontal line with intercept=0 (blue).

Predictor: s(albumin, 3)




Regression splines approach to determine non-linearity.
Variable vs. partial residual for smoothed variable.
If linear should be horizontal line with intercept=0 (blue)

Predictor: age with 4 df




Regression splines approach to determine non-linearity.
Variable vs. partial residual for smoothed variable.
If linear should be horizontal line with intercept=0 (blue)

Predictor: bili with 4 df




Regression splines approach to determine non-linearity.
Variable vs. partial residual for smoothed variable.
If linear should be horizontal line with intercept=0 (blue)

Predictor: protime with 4 df




Regression splines approach to determine non-linearity.
Variable vs. partial residual for smoothed variable.
If linear should be horizontal line with intercept=0 (blue)

Predictor: albumin with 4 df




Smoothing splines approach to determine non-linearity.
Variable vs. partial residual for smoothed variable.
If linear should be horizontal line with intercept=0 (blue)

Predictor: age with 4 df




Smoothing splines approach to determine non-linearity.
Variable vs. partial residual for smoothed variable.
If linear should be horizontal line with intercept=0 (blue)

Predictor: bili with 4 df




Smoothing splines approach to determine non-linearity.
Variable vs. partial residual for smoothed variable.
If linear should be horizontal line with intercept=0 (blue)

Predictor: protime with 4 df



Smoothing splines approach to determine non-linearity.
Variable vs. partial residual for smoothed variable.
If linear should be horizontal line with intercept=0 (blue)

Predictor: albumin with 4 df




Time vs. scaled Schoenfeld residuals, with smoothed spline (black).

If < 0 (red line), indicates a protective effect.

Regression line (blue) should be horizontal if model well fit.

Predictor: age
Time transform: km




Time vs. scaled Schoenfeld residuals, with smoothed spline (black).

If < 0 (red line), indicates a protective effect.

Regression line (blue) should be horizontal if model well fit.

Predictor: edema
Time transform: km




Time vs. scaled Schoenfeld residuals, with smoothed spline (black).

If <0 (red line), indicates a protective effect.

Regression line (blue) should be horizontal if model well fit.

Predictor: bili
Time transform: km




Time vs. scaled Schoenfeld residuals, with smoothed spline (black).

If < 0 (red line), indicates a protective effect.

Regression line (blue) should be horizontal if model well fit.

Predictor: protime
Time transform: km




Time vs. scaled Schoenfeld residuals, with smoothed spline (black).

If < 0 (red line), indicates a protective effect.

Regression line (blue) should be horizontal if model well fit.


Predictor: albumin

Time transform: km




Coefficient vs. jackknife influence.
Change in coefficient if this observation dropped.
Outliers may need to be re-examined

Coefficient: age




Coefficient vs. jackknife influence.
Change in coefficient if this observation dropped.
Outliers may need to be re-examined

Coefficient: edema




Coefficient vs. jackknife influence.
Change in coefficient if this observation dropped.
Outliers may need to be re-examined




Coefficient vs. jackknife influence.
Change in coefficient if this observation dropped.
Outliers may need to be re-examined

Coefficient: protime




Coefficient vs. jackknife influence.
Change in coefficient if this observation dropped.
Outliers may need to be re-examined

Coefficient: albumin




Coefficient vs. jackknife influence scaled by standard error of coefficients.
Change in coefficient if this observation dropped.
Outliers may need to be re-examined

Coefficient: age




Coefficient vs. jackknife influence scaled by standard error of coefficients.
Change in coefficient if this observation dropped.
Outliers may need to be re-examined

Coefficient: edema




Coefficient vs. jackknife influence scaled by standard error of coefficients.
Change in coefficient if this observation dropped.
Outliers may need to be re-examined

Coefficient: bili




Coefficient vs. jackknife influence scaled by standard error of coefficients.
Change in coefficient if this observation dropped.
Outliers may need to be re-examined

Coefficient: protime




Coefficient vs. jackknife influence scaled by standard error of coefficients.
Change in coefficient if this observation dropped.
Outliers may need to be re-examined

Coefficient: albumin




Coefficient vs. scaled score residuals.
Assesses leverage: influence of observation on a single coefficient.
Outliers may need to be re-examined

Coefficient: age




Coefficient vs. scaled score residuals.
Assesses leverage: influence of observation on a single coefficient.
Outliers may need to be re-examined

Coefficient: edema




Coefficient vs. scaled score residuals.
Assesses leverage: influence of observation on a single coefficient.
Outliers may need to be re-examined




Coefficient vs. scaled score residuals.
Assesses leverage: influence of observation on a single coefficient.
Outliers may need to be re-examined

Coefficient: protime




Coefficient vs. scaled score residuals.
Assesses leverage: influence of observation on a single coefficient.
Outliers may need to be re-examined

Coefficient: albumin




Martingale residuals vs. likelihood displacement residuals.
Assesses influence of observation on coefficient.
Outliers may need to be re-examined.

Coefficient: age




Martingale residuals vs. likelihood displacement residuals.
Assesses influence of observation on coefficient.
Outliers may need to be re-examined.

Coefficient: edema




Martingale residuals vs. likelihood displacement residuals.
Assesses influence of observation on coefficient.
Outliers may need to be re-examined.

Coefficient: bili



Martingale residuals vs. likelihood displacement residuals.
Assesses influence of observation on coefficient.
Outliers may need to be re-examined.

Coefficient: protime



Martingale residuals vs. likelihood displacement residuals.
Assesses influence of observation on coefficient.
Outliers may need to be re-examined.

Coefficient: albumin

