

# rgbif vignette - Search and retrieve data from the Global Biodiversity Information Facility (GBIF)

## About the package

rgbif is an R package to search and retrieve data from the Global Biodiversity Information Facility (GBIF). rgbif wraps R code around the [GBIF API](#) to allow you to talk to GBIF from R.

---

## Install rgbif and dependencies

```
install.packages("rgbif")
```

## Load rgbif

```
library(rgbif)
```

---

## Get number of occurrences for a set of search parameters

Search by type of record, all observational in this case

```
occ_count(basisOfRecord = "OBSERVATION")
```

```
[1] 100241780
```

## Records for Puma concolor with lat/long data (georeferenced) only

Note that hasCoordinate in occ\_search() is the same as georeferenced in occ\_count().

```
occ_count(taxonKey = 2435099, georeferenced = TRUE)
```

```
[1] 2604
```

## All georeferenced records in GBIF

```
occ_count(georeferenced = TRUE)
```

```
[1] 377067550
```

## Records from Denmark

```
occ_count(country = "DENMARK")
```

```
[1] 8645654
```

## Number of records in a particular dataset

```
occ_count(datasetKey = "9e7ea106-0bf8-4087-bb61-dfe4f29e0f17")
```

```
[1] 4591
```

## All records from 2012

```
occ_count(year = 2012)
```

```
[1] 32234055
```

## Records for a particular dataset, and only for preserved specimens

```
occ_count(datasetKey = "e707e6da-e143-445d-b41d-529c4a777e8b", basisOfRecord = "OBSERVATION")
```

```
[1] 2120907
```

---

## Get possible values to be used in taxonomic rank arguments in functions

```
taxrank()
```

```
[1] "kingdom"    "phylum"   "class"      "order"
[5] "family"     "genus"      "species"    "infraspecific"
```

---

## Search for taxon information

### Search for a genus

```
head(name_lookup(query = "Cnaemidophorus", rank = "genus", return = "data"))
```

	key	nubKey	parentKey	parent	kingdom	phylum	clazz
1	116755723	1858636	110614854	Pterophoridae	Animalia	Arthropoda	Insecta
2	1858636	1858636	8863	Pterophoridae	Animalia	Arthropoda	Insecta
3	125802004	1858636	125793784	Pterophoridae	<NA>	<NA>	Insecta
4	131295800	1858636	NA	<NA>	<NA>	<NA>	<NA>
5	127882857	1858636	127804516	Pterophoridae	Animalia	Arthropoda	Insecta
6	115123697	1858636	NA	<NA>	<NA>	<NA>	<NA>

	order	family	genus	kingdomKey	phylumKey	classKey
1	Lepidoptera	Pterophoridae	Cnaemidophorus	116630539	116762374	131743724
2	Lepidoptera	Pterophoridae	Cnaemidophorus	1	54	216
3	Lepidoptera	Pterophoridae	Cnaemidophorus	NA	NA	131714461
4	<NA>	<NA>	Cnaemidophorus	NA	NA	NA
5	Lepidoptera	Pterophoridae	Cnaemidophorus	127795487	127795488	127795683
6	<NA>	<NA>	Cnaemidophorus	NA	NA	NA

	orderKey	familyKey	genusKey	canonicalName	authorship	nameType
1	116843281	110614854	116755723	Cnaemidophorus	Wallengren, 1862	WELLFORMED
2	797	8863	1858636	Cnaemidophorus	Wallengren, 1862	WELLFORMED
3	125810165	125793784	125802004	Cnaemidophorus	Wallengren, 1862	WELLFORMED
4	NA	NA	131295800	Cnaemidophorus		WELLFORMED
5	127795981	127804516	127882857	Cnaemidophorus		WELLFORMED
6	NA	NA	115123697	Cnaemidophorus		WELLFORMED

	rank	numOccurrences
1	GENUS	0
2	GENUS	0
3	GENUS	0
4	GENUS	0
5	GENUS	0
6	GENUS	0

Search for the class mammalia

```
head(name_lookup(query = "mammalia")$data)
```

	key	nubKey	parentKey	parent	phylum	clazz
1	125798198	359	131712102	Chordata	Chordata	Mammalia
2	116665331	359	116842680	Chordata	Chordata	Mammalia
3	359	359	44	Chordata	Chordata	Mammalia
4	125826646	359	131712102	Chordata	Chordata	Mammalia
5	131754503	359	131754502	Macroscelidea	Chordata	Macroscelidea
6	102402290	359	102545028	Chordata	Chordata	Mammalia

	phylumKey	classKey	canonicalName	authorship	nameType	rank
1	131712102	125798198	Mammalia		WELLFORMED	CLASS
2	116842680	116665331	Mammalia	Linnaeus, 1758	WELLFORMED	CLASS
3	44	359	Mammalia	Linnaeus, 1758	WELLFORMED	CLASS
4	131712102	125826646	Mammalia	Linnaeus, 1758	WELLFORMED	CLASS
5	116842680	131754502	Mammalia		WELLFORMED	ORDER
6	102545028	102402290	Mammalia		WELLFORMED	CLASS

	numOccurrences	kingdom	kingdomKey	order	orderKey
--	----------------	---------	------------	-------	----------

1	0	<NA>	NA	<NA>	NA
2	0	Animalia	116630539	<NA>	NA
3	0	Animalia	1	<NA>	NA
4	0	<NA>	NA	<NA>	NA
5	0	Animalia	116630539	Mammalia	131754503
6	0	Animalia	101719444	<NA>	NA

Look up the species *Helianthus annuus*

```
head(name_lookup("Helianthus annuus", rank = "species")$data)
```

	key	nubKey	parentKey	parent	kingdom	order
1	116845199	3119195	116853573	Helianthus	Plantae	Asterales
2	3119195	3119195	3119134	Helianthus	Plantae	Asterales
3	125790787	3119195	125809269	Helianthus	<NA>	Asterales
4	106239436	3119195	106239325	Helianthus	Viridiplantae	Asterales
5	128399814	3119195	131338207	Helianthus	<NA>	<NA>
6	111449704	3119195	111449703	Helianthus	Plantae	<NA>
	family	genus	kingdomKey	orderKey	familyKey	genusKey
1	Asteraceae	Helianthus	116668764	116852024	116856030	116853573
2	Asteraceae	Helianthus	6	414	3065	3119134
3	Asteraceae	Helianthus	NA	131717243	125799038	125809269
4	Asteraceae	Helianthus	106147210	106237428	106237535	106239325
5	<NA>	Helianthus	NA	NA	NA	131338207
6	Compositae	Helianthus	111449174	NA	111442813	111449703
	canonicalName	authorship	nameType	rank	numOccurrences	
1	Helianthus annuus	L.	WELLFORMED	SPECIES	0	
2	Helianthus annuus	L.	WELLFORMED	SPECIES	0	
3	Helianthus annuus	L.	WELLFORMED	SPECIES	0	
4	Helianthus annuus		WELLFORMED	SPECIES	0	
5	Helianthus annuus		WELLFORMED	SPECIES	0	
6	Helianthus annuus	L.	WELLFORMED	SPECIES	0	
	phylum	clazz	phylumKey	classKey		
1	<NA>	<NA>	NA	NA		
2	Magnoliophyta	Magnoliopsida	49	220		
3	<NA>	<NA>	NA	NA		
4	Streptophyta	<NA>	106171079	NA		
5	<NA>	<NA>	NA	NA		
6	Spermatophyta	Dicotyledones	111449175	111449177		

Get data for a single occurrence. Note that data is returned as a list, with slots for metadata and data, or as a hierarchy, or just data.

Just data

```
occ_get(key = 766766824, return = "data")
```

	name	key	decimalLatitude	decimalLongitude
1	Corvus monedula	766766824	59.46	17.91

## Just taxonomic hierarchy

```
occ_get(key = 766766824, return = "hier")
```

	name	key	rank
1	Animalia	1	kingdom
2	Chordata	44	phylum
3	Aves	212	class
4	Passeriformes	729	order
5	Corvidae	5235	family
6	Corvus	2482468	genus
7	Corvus monedula	2482473	species

## All data, or leave return parameter blank

```
occ_get(key = 766766824, return = "all")
```

\$hierarchy

	name	key	rank
1	Animalia	1	kingdom
2	Chordata	44	phylum
3	Aves	212	class
4	Passeriformes	729	order
5	Corvidae	5235	family
6	Corvus	2482468	genus
7	Corvus monedula	2482473	species

\$data

	name	key	decimalLatitude	decimalLongitude
1	Corvus monedula	766766824	59.46	17.91

## Get many occurrences. occ\_get is vectorized

```
occ_get(key = c(766766824, 101010, 240713150, 855998194, 49819470), return = "data")
```

	name	key	decimalLatitude	decimalLongitude
1	Corvus monedula	766766824	59.46	17.91
2	Platydoras costatus	101010	-4.35	-70.07
3	<NA>	240713150	-77.57	163.58
4	Sciurus vulgaris	855998194	58.41	12.04
5	Phlogophora meticulosa	49819470	55.72	13.28

## Maps

### Static map using the ggplot2 package

Make a map of **Puma concolor** occurrences

```
key <- name_backbone(name = "Puma concolor", kingdom = "plants")$speciesKey
dat <- occ_search(taxonKey = key, return = "data", limit = 300)
gbifmap(input = dat)
```




Figure 1: plot of chunk gbifmap1